

## Risks associated with your anaesthetic

# SECTION 5: DAMAGE TO THE EYE DURING GENERAL ANAESTHESIA

**During a general anaesthetic it is possible for your eyes to be damaged. This is an uncommon or rare event. The types of damage that can occur, and its consequences and treatment are described in this article.**

**Damage caused during surgery to the eye itself or associated with anaesthesia for eye surgery is not described here. You can talk about this with the eye surgeon or specialist anaesthetist who is looking after you.**

### What is the most common type of damage?

The most common damage to the eye which can occur during or after general anaesthesia is a corneal abrasion.<sup>1,2</sup>

### What is a corneal abrasion?

The cornea is one of the outer clear layers of the eye. An abrasion is a tear or graze of this layer. The abrasion can cause pain, blurred vision and considerable irritation for a few days. Almost all corneal abrasions heal without long-term effects on vision.

### How do corneal abrasions happen?

Most abrasions happen because the eye does not close fully during the anaesthetic. Approximately 6 out of 10 people (60%) do not close their eyes naturally when they have a general anaesthetic.<sup>1</sup> The cornea is then exposed to the air and becomes dry. Fewer tears are produced during an anaesthetic which also causes dryness in the eyes.<sup>2,3</sup>

The dry cornea can then stick to the inside of the eyelid and the abrasion occurs when the eye opens again at the end of the anaesthetic.

Corneal abrasion can also occur because something rubs against the exposed cornea during the anaesthetic. This may be one of the sheets used during surgery to cover the patient and keep the operation area sterile, or other equipment. Anaesthetists take care to ensure the eyes are closed during a general anaesthetic and to protect the eyes.

### What is done to prevent corneal abrasions?

Corneal abrasions can usually be prevented by careful protection of the eyes.

Small pieces of sticking tape are commonly used to keep the eyelids fully closed during the anaesthetic. This has been shown to reduce the chance of a corneal abrasion occurring.<sup>1,2</sup> However, bruising of the eyelid can occur when the tape is removed, especially if you have thin skin and bruise easily.

Sometimes, the anaesthetist may use a gel, an ointment or eye drops to moisten the eyes during your anaesthetic. These may be helpful if tape cannot be used or for certain operations in which the eyes need to be opened briefly during the operation. Eye ointments can

sometimes cause temporary eye irritation or blurring of vision following an anaesthetic.

Anaesthetists are trained to take care that nothing rubs against the eyes. If your surgery requires you to be positioned lying on your front, your anaesthetist will use goggles, cushions and/or eyepads to protect your eyes.

### How often do corneal abrasions occur?

Following a general anaesthetic, it is uncommon to suffer from a corneal abrasion that causes symptoms. A large study of over 60,000 patients having a general anaesthetic found that 1 in 2,800 patients suffered symptoms from a corneal abrasion.<sup>4</sup>

Studies have also been done using a microscope to examine the eyes following an anaesthetic. These show that small corneal abrasions occur commonly. Around 1 in 25 patients may have a small corneal abrasion, which the patient does not notice. This occurs even when protective eye tape or ointment is used.

You are more likely to suffer from a corneal abrasion if your surgery requires you to be positioned lying on your front or your side, if your operation lasts a long time, or if you are having surgery on your head or neck.<sup>4</sup>

### What if I already have poor vision?

If you have poor vision, it is helpful if you tell your anaesthetist about it. This is because he/she can give you any extra information that you need to help you feel at ease if you cannot see well. However, this will not make any difference to the risk of getting a corneal abrasion, or to the ways in which your anaesthetist cares for your eyes while you are anaesthetised.

### What happens if I have a corneal abrasion?

Corneal abrasions may be very painful. Healing usually takes a few days, after which the pain will stop completely. Treatment during this time can reduce pain and aims to prevent an eye infection developing. Eye drops, ointments and an eye patch may be used, as well as pain-relieving medicines. No surgical treatment is necessary.

Almost all corneal abrasions heal with no visible scar and no long-term effect on vision. An eye specialist may be able to see a scar through a microscope. Contact lens users should take advice before using contact lenses again.

### Can I lose my sight during a general anaesthetic?

Serious eye injuries during a general anaesthetic are very rare, but can lead to loss of eyesight. Two structures can be damaged:

- ▶ the retina is the light-sensitive layer inside the eye
- ▶ the optic nerve carries visual information from the eye to the brain.

One or both of these can be damaged by one or more of the following:

- ▶ low blood pressure during the operation can mean that the optic nerve and/or the retina do not get enough oxygen
- ▶ tiny clots in the blood vessels to the eye can also cause lack of oxygen to the optic nerve or the retina
- ▶ too much pressure on the eyeball during the operation can damage the optic nerve or interrupt the blood supply to the nerve and the retina.

The operations with higher risk are:

- ▶ operations on the spine in the prone position (face down), especially if the operation lasts more than six hours and there is a lot of loss of blood<sup>5</sup>
- ▶ operations which require cardio-pulmonary bypass (open heart surgery with use of a heart/lung machine)
- ▶ neck dissection operations on both sides of the neck.

During these operations the anaesthetist will take particular care to keep the blood pressure at an appropriate level and prevent pressure on the eyeball.

The people with higher risk include those with other vascular diseases (high blood pressure, heart attack or stroke), diabetics and those with high red blood cell counts.

Overall, it is very rare to lose sight in an eye after a general anaesthetic. It is difficult to give an accurate figure for the risk because it is so rare. Studies suggest that the overall risk, for all operations under general anaesthetic is between 1 in 60,000 and 1 in 125,000 operations.<sup>2,3,6,7</sup> However it is more likely (but still uncommon) in the high risk operations listed above. One study estimates that visual loss happens in 1 in 3,300 operations on the spine and 1 in 1,100 open heart operations.<sup>7</sup>

## Are there any other eye injuries that can happen?

Other eye problems that can follow a general anaesthetic include:

- ▶ Pressure on nerves in the eyebrow area may cause a droopy eyelid. This is usually temporary and should recover.
- ▶ Protective tape or eye ointments used to protect your eyes from corneal abrasions may cause temporary bruising of the eyelids or irritation of the eyes. Redness of the eye, blurred vision and the feeling that there is something in the eye may last for up to eight hours.
- ▶ A few operations are performed in an extreme head down position. This includes prolonged laparoscopic (keyhole) surgery for major procedures such as removal of parts of the bowel. This can lead to swelling of your eyelids which should resolve within a short time.
- ▶ If you have glaucoma (which causes high pressure inside your eye) your anaesthetist will need to take extra care to protect your eyesight during surgery.

### Summary

Corneal abrasion causing symptoms is an uncommon eye problem after general anaesthesia. Anaesthetists take care to prevent it by using tape or other forms of eye protection. It is more common in patients who need to lie face down for surgery.

Loss of sight after general anaesthesia is very rare, but should be considered a risk in certain operations.

### Author (2009)

**Dr Lucy White**, FRCA  
Consultant Anaesthetist  
University Hospital Southampton  
NHS Foundation Trust

### Reviewer 2013 edition

**Dr Andrew Presland**, FRCA  
Consultant Anaesthetist  
Moorfields Eye Hospital

### References

1. White E, Crosse MM. The aetiology and prevention of peri-operative corneal abrasions. *Anaesthesia* 1998;**53**:157–161.
2. White E. Care of the eye during anaesthesia. *Anaesth Intensive Care* 2004;**5**:302–303.
3. Contractor S, Hardman JG. Injury during anaesthesia. Continuing Education in Anaesthesia, Critical care and Pain. Volume 6 number 2 2006.
4. Roth S, Thisted RA, Erickson JP et al. Eye injuries after non ocular surgery: a study of 60,965 anaesthetics from 1988 to 1992. *Anesthesiology* 1996;**85**:1020–1027.
5. Lee LA et al. The American Society of Anesthesiologists Postoperative Visual Loss Registry: analysis of 93 spine surgery cases with postoperative visual loss. *Anesthesiology* 2006;**105**(4): 652–659.
6. Rupp-Montpetit K, Moody ML. Visual loss as a complication of non-ophthalmic surgery: a review of the literature. *AANA Journal* 2004;**72**(4):285–292.
7. Roth S. Peri-operative visual loss : what do we know, what can we do? *Br J Anaesth* 2009;**103**:i31–i40.



The Royal College of  
Anaesthetists

### The Royal College of Anaesthetists

Revised edition 2013

The material from this article may be copied for the purpose of producing information materials for patients. Please quote the RCoA as the source of the information. If you wish to use part of the article in another publication, suitable acknowledgement must be given and the RCoA logo must be removed. For more information or enquiries about the use of this leaflet please contact:

The Royal College of Anaesthetists

website: [www.rcoa.ac.uk](http://www.rcoa.ac.uk)

email: [comms@rcoa.ac.uk](mailto:comms@rcoa.ac.uk)

This leaflet will be reviewed three years from the date of publication.